

Post-stroke mRNA expression profile of MMPs: effect of genetic deletion of MMP-12

Koteswara Rao Nalamolu,¹ Bharath Chelluboina,¹ Ian B Magruder,¹ Diane N Fru,¹ Adithya Mohandass,¹ Ishwarya Venkatesh,¹ Jeffrey D Klopfenstein,^{1,2,3} David M Pinson,⁴ Krishna M Boini,⁵ Krishna Kumar Veeravalli^{1,2,6}

To cite: Nalamolu KR, Chelluboina B, Magruder IB, *et al.* Post-stroke mRNA expression profile of MMPs: effect of genetic deletion of MMP-12. *Stroke and Vascular Neurology* 2018;**0**: e000142. doi:10.1136/svn-2018-000142

Received 6 January 2018
Revised 7 February 2018
Accepted 7 February 2018



¹Department of Cancer Biology and Pharmacology, University of Illinois College of Medicine, Peoria, Illinois, USA

²Department of Neurosurgery, University of Illinois College of Medicine, Peoria, Illinois, USA

³Comprehensive Stroke Center, Illinois Neurological Institute, OSF HealthCare System, Saint Francis Medical Center, Peoria, Illinois, USA

⁴Department of Pathology, University of Illinois College of Medicine, Peoria, Illinois, USA

⁵Department of Pharmacological and Pharmaceutical Sciences, College of Pharmacy, University of Houston, Houston, Texas, USA

⁶Department of Neurology, University of Illinois College of Medicine, Peoria, Illinois, USA

Correspondence to

Dr Krishna Kumar Veeravalli; krishnav@uic.edu

ABSTRACT

Background and purpose Recent reports from our laboratory demonstrated the post-ischaemic expression profile of various matrix metalloproteinases (MMPs) in rats and the detrimental role of MMP-12 in post-stroke brain damage. We hypothesise that the post-stroke dysregulation of MMPs is similar across species and that genetic deletion of MMP-12 would not affect the post-stroke expression of other MMPs. We tested our hypothesis by determining the pre-ischaemic and post-ischaemic expression profile of MMPs in wild-type and MMP-12 knockout mice.

Methods Focal cerebral ischaemia was induced in wild-type and MMP-12 knockout mice by middle cerebral artery occlusion procedure by insertion of a monofilament suture. One hour after ischaemia, reperfusion was initiated by removing the monofilament. One day after reperfusion, ischaemic brain tissues from various groups of mice were collected, and total RNA was isolated and subjected to cDNA synthesis followed by PCR analysis.

Results Although the post-stroke expression profile of MMPs in the ischaemic brain of mice is different from rats, there is a clear species similarity in the expression of MMP-12, which was found to be predominantly upregulated in both species. Further, the post-stroke induction or inhibition of various MMPs in MMP-12 knockout mice is different from their respective expression profile in wild-type mice. Moreover, the brain mRNA expression profile of various MMPs in MMP-12 knockout mice under normal conditions is also different to their expression in wild-type mice.

Conclusions In the ischaemic brain, MMP-12 upregulates several fold higher than any other MMP. Mice derived with the genetic deletion of MMP-12 are constitutive and have altered MMP expression profile both under normal and ischaemic conditions.

INTRODUCTION

The post-stroke dysregulation of matrix metalloproteinases (MMPs) during the acute phases after ischaemia and reperfusion has been associated with increased neurovascular disruption and brain injury. Post-stroke dysregulation of MMPs leads to blood–brain barrier disruption, cerebral oedema, haemorrhage, leucocyte infiltration and inflammation.^{1–6} In the early stages after ischaemic

stroke, MMPs participate in the injury process. However, during the later stages after stroke, MMPs contribute to recovery. MMP inhibitors have shown promise in animal models of ischaemic stroke. However, inhibition of MMPs during the recovery phase after stroke increased brain injury, suppressed neurovascular remodelling and impaired functional recovery.⁷ Therefore, the timing of the inhibition of MMPs after stroke is critical. It is the timing of MMP inhibition and also the type of MMP that is inhibited that influences the overall outcome. Upregulation of MMP-1, MMP-2, MMP-3, MMP-8, MMP-9, MMP-10 and MMP-13 was reported in human ischaemic brain tissues.⁸ Of all the MMPs, MMP-2 and MMP-9 were well reported for their role in the context of ischaemic brain damage. Recent literature highlighted both detrimental and beneficial roles of several other MMPs such as MMP-3, MMP-8, MMP-10, MMP-12 and MMP-13 in the context of ischaemic stroke.^{6,9–15}

Because of the critical role various MMPs have on injury and during the recovery process after ischaemic stroke, investigating the post-stroke expression profile of all the known MMPs and understanding their contribution to the injury/recovery process would help the researchers to discover novel strategies to treat ischaemic stroke. We recently reported in a rat model the temporal expression profile of the majority of the known MMPs at various reperfusion times subsequent to a 2-hour focal cerebral ischaemia.¹⁰ We hypothesise that the post-stroke dysregulation of MMPs is similar across species. In this study, we tested our hypothesis by evaluating the post-stroke expression profile of these MMPs at 1-day reperfusion in a mouse model of ischaemic stroke. We also hypothesise that the genetic deletion of any MMP, for example MMP-12 in this study, would not affect the post-stroke

expression of other MMPs. We tested our hypothesis by determining the pre-ischaemic and post-ischaemic expression profile of MMPs in wild-type as well as MMP-12 knockout mice. Our results indicated a predominant upregulation of MMP-12 in the ischaemic brain of mice. The degree of post-stroke induction of MMP-12 as compared with other MMPs in mice is in agreement with our recent findings in rats. The post-stroke expression of other MMPs except MMP-9 is different in mice and rats. Further, the genetic deletion of MMP-12 altered both the pre-ischaemic and post-ischaemic expression of MMPs compared with wild-type mice.

METHODS

Ethics statement

The Institutional Animal Care and Use Committee (IACUC) of the University of Illinois College of Medicine at Peoria approved the study design, surgical manipulations, postoperative care and humane endpoints. All the procedures performed on animals complied with the approved IACUC protocol.

Animals and experimental design

Healthy, young male MMP-12 knockout (strain: B6.129X-Mmp12^{tm1Sds}/J) and respective wild-type (strain: C57BL/6J) mice procured from the Jackson Laboratory (USA) were used in this study. Animals were housed in a 12-hour light/dark cycle with a controlled temperature and humidity and free access to food and water. All animal experiments were conducted in accordance with the Guide for the Care and Use of Laboratory Animals and the approved IACUC protocol. Mice were randomised to various groups. Middle cerebral artery occlusion (MCAO) subjected wild-type and MMP-12 knockout mice were from groups 2 and 4, respectively. Sham-operated wild-type and MMP-12 knockout mice were from groups 1 and 3, respectively. All animals from the four groups were euthanised 1 day post-MCAO procedure. Because of the mortality associated with the MCAO procedure, we increased the number of animals in each group to attain the required sample size for statistical analysis. Animals that did not show post-stroke symptoms after MCAO procedure were excluded from the study.

Induction of focal cerebral ischaemia and reperfusion

After the animals reached a weight of ~23.5g, they were subjected to right MCAO procedure by using a silicone rubber-coated monofilament suture (Doccol, California, USA). Briefly, a ventral midline incision (~15–20mm) was made in the neck and the right common carotid, internal carotid and external carotid arteries were surgically exposed. The external carotid artery (ECA) was permanently ligated rostral with one ligature. Another loose ligature was made to the ECA near the bifurcation. Two microaneurysm clips were applied each to the common carotid artery (CCA) and internal carotid artery (ICA). A small puncture opening was made to the ECA between the two ligatures. The monofilament

was inserted through the opening on ECA and advanced into ICA. The microaneurysm clip was removed from ICA, and the monofilament was gently further advanced into ICA up to 9 to 10mm distance that is pre-marked on the monofilament. The other loose ligature was tightened around the ECA containing the monofilament. The microaneurysm clip on CCA was removed, and skin on the neck incision was closed with surgical wound clips. To restore the blood flow 1 hour after MCAO, the surgical site was reopened by removing the wound clips. The microaneurysm clip was again applied to CCA, the knot was loosened, the monofilament was withdrawn and the knot was re-tied to stop bleeding. The microaneurysm clip on CCA was removed and skin was sutured to close the neck incision.

RNA extraction, cDNA synthesis and real-time PCR analysis

From the entire ipsilateral hemisphere of mice brains, total RNA was extracted using TRIzol reagent (Invitrogen, USA), reverse transcribed to cDNA with iScript cDNA synthesis kit (Bio-Rad Laboratories, USA) and used for real-time PCR analysis by using SYBR Green method. Briefly, reaction set-up for each cDNA sample was assembled using the FastStart SYBR Green Master (Roche, Indianapolis, Indiana, USA) as per the manufacturer's instructions. Samples were subjected to 40 cycles at 95°C for 15 s, 60°C for 1 min and 72°C for 1 min in iCycler IQ (Multi Colour Real-Time PCR Detection System; Bio-Rad Laboratories, Hercules, California, USA). Data were collected and recorded using the iCycler IQ software (Bio-Rad Laboratories) and expressed as a function of the threshold cycle (Ct), which represents the number of cycles at which the fluorescent intensity of the SYBR Green dye is significantly above that of the background fluorescence. The genes and their respective forward and reverse primer sequences analysed by real-time PCR are listed in [table 1](#). β -Actin served as a housekeeping gene. Average of Ct values were normalised with average Ct values of β -actin. After normalisation of the Ct values, fold differences were calculated by using the formula $2^{-(\Delta C_t \text{ of test})} / 2^{-(\Delta C_t \text{ of controls})}$.

RT-PCR analysis and agarose gel electrophoresis

To validate the changes in mRNA expression among groups, RT-PCR analysis was performed for certain genes such as MMP-12 and MMP-9 along with the housekeeping gene β -actin using GoTaq Green Master Mix (Promega) as per the manufacturer's instructions. RT-PCR was set up in C1000 Touch Thermo cycler (Bio-Rad Laboratories) using the following PCR cycle: (95°C for 5 min (95°C for 30 s, 58°C–62°C for 45 s and 72°C for 45 s) \times 35 cycles, and 72°C for 10 min). RT-PCR products were resolved on 2% agarose gels containing ethidium bromide, visualised and photographed under UV light.

Statistical analysis

Statistical analysis of the data was performed using GraphPad Prism software (V.3.02). Quantitated data of

Table 1 Mouse-specific genes analysed by PCR

Gene	Accession number	Forward primer (5'–3')	Reverse primer (5'–3')
MMP-1a	NM_032006	ccttccttctgtctgcttc	ctccttgccattcacgttt
MMP-1b	NM_032007	gtgaatggcaaggaggtgat	ggccaacgaggattgtgt
MMP-2	NM_008610	gtcgcccctaaaacagacaa	ggctcgtatggtgtctggt
MMP-3	NM_010809	cagactgtcccgtttccat	ggtgtgactgcatcaaaga
MMP-7	NM_010810	cggagatgctcactttgaca	cagcgtgttctctttccat
MMP-8	NM_008611	gcctcccagtagctgaaca	actccacatcgaggcatttc
MMP-9	NM_013599	cgctgtatccccacttact	aacacacagggtttgccttc
MMP-10	NM_019471	gacccactcactttctcca	gggtgcaagtgtccatttct
MMP-11	NM_008606	ccctccaggtatggagtga	caggtcagttccctggtgt
MMP-12	M82831	ccaagcatcccatctgctat	ggtaaagacagctgcatca
MMP-13	NM_008607	tttattgttctgcccata	ctctggtgtttgggatgct
MMP-14	NM_008608	ccgcatgcaaaagtctat	gccacccttaggggtgtaat
MMP-15	NM_008609	cccacaggtcacaccttct	ccagttactggtgcccctgt
MMP-16	NM_019724	ggagacagttcccattga	ccagctcatggactgtaca
MMP-17	NM_011846	caccactttgatgacgatg	ccctggtagtacgggtgcat
MMP-21	NM_152944	gactgggcagacagaaaagc	gagtcctgtgttgccggtt
MMP-25	NM_001033339	gctgactcgctatggctacc	cttccaacactgccactca
MMP-28	NM_001320300	gaggcgtaagaacgctttg	ccagaactccagtgctgaca
βactin	NM_007393.5	agccatgtacgtagccatcc	ctctcagctgtggtggtgaa

PCR analysis were tested for normality and equality of variances between groups. The data that were normally distributed and have equal variances between groups were

Table 2 Matrix metalloproteinases (MMPs) that are either upregulated or downregulated after focal cerebral ischaemia and reperfusion in rats and mice

Gene	Δ Fold over sham in rats	Δ Fold over sham in mice
MMP-1a	No change	–13.08
MMP-2	No change	11.12
MMP-3	11.28	No change
MMP-8	No change	279.55
MMP-9	16.05	24.39
MMP-10	No change	–247.83
MMP-12	47.15	763.79
MMP-15	No change	–22.33
MMP-16	No change	–29.74
MMP-17	No change	–32.47
MMP-21	No change	–53.60

Focal cerebral ischaemia was induced for 2 hours in rats and 1 hour in mice. Changes in mRNA expression of various MMPs in the ischaemic brain of rats versus mice were analysed 1 day after reperfusion. MMPs that were either upregulated or downregulated significantly ($P < 0.05$) and more than 10-fold as compared with their respective sham-operated rats¹⁰ and mice are included in this table and those with less than 10-fold change are indicated as 'no change'. A positive value indicates upregulation and a negative value indicates downregulation. $n = 6$.

tested for differences by unpaired t-test. We applied Welch's correction to unpaired t-test for the data that were normally distributed but has unequal variances between groups. Results are expressed as the mean \pm SD. Differences in the values were considered significant at $P < 0.05$.

RESULTS

Dysregulation of MMPs after ischaemic stroke is not the same in rats and mice

Our recent study demonstrated an upregulation (>10 -fold vs sham) of MMP-3, MMP-9 and MMP-12 in ischaemic rat brains at 1 day reperfusion after a 2-hour focal cerebral ischaemia.¹⁰ Further, we noticed that the post-stroke induction of MMP-12 in rats was higher than any other MMPs tested. In contrast to the expression pattern that was previously reported in the ischaemic brain of rats, in this study, MMP-2, MMP-8, MMP-9 and MMP-12 were upregulated (>10 -fold vs sham) and MMP-1a, MMP-10, MMP-15, MMP-16, MMP-17 and MMP-21 were downregulated (>10 -fold vs sham) in the ischaemic brains of mice (table 2). Of all the MMPs tested, MMP-12 upregulation (~ 763 -fold vs sham) was predominant in mice. Post-stroke induction of MMP-8 (~ 279 -fold vs sham) in mice was the largest after MMP-12. Interestingly, marked downregulation of MMP-10 (~ 247 -fold vs sham) expression was noticed in mice.

Genetic deletion of MMP-12 alters the post-stroke dysregulation of MMPs

There is no difference in the post-stroke expression of MMP-12 in the ischaemic brain of MMP-12 knockout mice

as compared with their sham. As expected, significant upregulation in the post-stroke expression of MMP-12 was noticed in wild-type mice (figure 1A; unpaired t-test with Welch's correction; Welch corrected $t=3.608$; $P=0.0154$, $n=6$). Genetic deletion of MMP-12 did not affect the post-stroke expression of MMP-9 (figure 1A, B). As mentioned earlier, marked upregulation of MMP-8 (~279-fold vs sham) and downregulation of MMP-10 (~247-fold vs sham) were noticed in wild-type mice subjected to 1-hour ischaemia followed by 1-day reperfusion (figure 2).

Genetic deletion of MMP-12 in mice showed a significant difference as compared with wild-type mice in the post-stroke expression of several MMPs such as MMP-1a (figure 2; unpaired t-test with Welch's correction; Welch corrected $t=3.699$; $P=0.0140$, $n=6$), MMP-2 (figure 2; unpaired t-test with Welch's correction; Welch corrected $t=5.062$; $P=0.0023$, $n=6$), MMP-8 (figure 2; unpaired t-test with Welch's correction; Welch corrected $t=4.122$; $P=0.0092$, $n=6$), MMP-10 (figure 2; unpaired t-test with Welch's correction; Welch corrected $t=3.812$; $P=0.0125$, $n=6$), MMP-21 (figure 2; unpaired t-test with Welch's correction; Welch corrected $t=5.930$; $P=0.0019$, $n=6$) and MMP-28 (figure 2; unpaired t-test with Welch's correction; Welch corrected $t=4.773$; $P=0.005$, $n=6$). As mentioned earlier, MMP-15, MMP-16 and MMP-17 were downregulated (>10-fold vs sham) in the ischaemic brain of wild-type mice (table 2). Although the downregulation of MMP-15, MMP-16 and MMP-17 was prevented in MMP-12 knockout mice as compared with wild-type mice, the differences were not significant.

Expression pattern of MMPs is not similar in MMP-12 knockout and wild-type mice

As we noticed significant differences in the expression pattern of various MMPs in the ischaemic brain of wild-type and MMP-12 knockout mice that were subjected to 1-day reperfusion after 1-hour focal cerebral ischaemia, we tested the mRNA expression pattern of MMPs in these mice and under normal conditions. Our results showed that the expression of MMPs in MMP-12 knockout mice was different to their expression in wild-type mice. Except MMP-8, which showed an increased expression (~26-fold vs wild type), the expression of almost all the MMPs tested was less in MMP-12 knockout mice as compared to their expression level in wild-type mice under normal conditions (figure 3). Interestingly, the decrease in the mRNA expression of MMP-10, MMP-15, MMP-21 and MMP-28 was prominent (>50-fold vs wild type) and the decrease was more than the degree of reduction in MMP-12 mRNA expression in MMP-12 knockout mice.

DISCUSSION

In this study, we showed the post-stroke expression profile of various MMPs in mouse brain after focal cerebral ischaemia and reperfusion. We noticed that the induction or inhibition of MMPs in mice after ischaemic stroke was different to the expression pattern that we recently




Figure 1 Post-stroke expression of MMP-12 and MMP-9 in the ischaemic brain of wild-type and MMP-12 knockout mice. (A) Real-time PCR analysis of samples from the ischaemic brains of mice at 1 day after reperfusion subsequent to a 1-hour focal cerebral ischaemia. Histograms and error bars indicate the mean and the SD, respectively. $n=6$. * $P<0.05$ versus wild type. (B) Bands represent the mRNA expression of MMP-12 and MMP-9. Beta actin served as a loading control. I, ischaemia; M12KO, MMP-12 knockout mice; R, reperfusion.

reported in rats.¹⁰ While comparing the expression pattern of MMPs in mice versus rats, we here discuss only the increasing or decreasing trend but not the degree of increase or decrease, as the induction time of ischaemia (1 hour in mice vs 2 hours in rats) in these rodent models is different. MMP-3, MMP-9 and MMP-12 were upregulated (>10-fold vs sham) and none of the MMPs tested were downregulated in rats.¹⁰ In contrast to rats, in this study, MMP-2, MMP-8, MMP-9 and MMP-12 were upregulated (>10-fold vs sham), whereas MMP-1a, MMP-10,




Figure 2 Dysregulation of MMPs in the ischaemic brain of wild-type and MMP-12 knockout mice. Real-time PCR analysis of samples from the ischaemic brains of mice at 1 day after reperfusion subsequent to a 1-hour focal cerebral ischaemia. Histograms and error bars indicate the mean and the SD, respectively, $n=6$. * $P<0.05$ vs wild type, ** $P<0.01$ versus wild type. M12KO, MMP-12 knockout mice; WT, wild-type mice.

MMP-15, MMP-16, MMP-17 and MMP-21 were down-regulated (>10 -fold vs sham) in mice. MMP-1, MMP-2, MMP-3, MMP-8, MMP-9, MMP-10 and MMP-13 proteins were upregulated in the ischaemic brain tissues of deceased patients who had an ischaemic stroke within the previous 4 days.⁸ The degree of severity of ischaemia and the presence or absence of reperfusion could be the possible reasons for the observed differences in the expression of MMPs in humans and rodent species.

Despite the differences in the expression pattern of various MMPs in mice versus rats, MMP-9 was upregulated in the ischaemic brain of both these rodent species at 1-day reperfusion. Consistent with our findings, in a recent review, authors have listed the studies that showed the induction of MMP-9 in these animal models and suggested the inhibition of MMP-9 (either by chemical inhibitors, MMP-9 neutralising antibodies or MMP-9 specific siRNAs/shRNAs) as a therapeutic strategy to attenuate post-stroke brain damage.¹⁶ The post-stroke induction of MMP-9 was also reported in non-human primates

and in human brain.^{8 17–19} It was shown in different ischaemic models that the genetic deletion of MMP-9 in mice attenuates the post-stroke brain damage.^{20–22}

In this study, MMP-12 expression was predominantly upregulated (~ 763 -fold vs sham) in mouse ischaemic brains. The post-stroke induction of MMP-12 in mice is consistent with our earlier findings in rats, wherein we reported a predominant upregulation of MMP-12 (~ 47 -fold vs sham at 1-day reperfusion and ~ 265 -fold vs sham at 7-day reperfusion) over other MMPs.¹⁰ MMP-12 can activate other MMPs such as pro-MMP-2 and pro-MMP-3, which, in turn, can activate pro-MMP-1 and pro-MMP-9.²³ Our recent reports clearly demonstrated the detrimental role of the induced MMP-12 on post-stroke brain damage.^{6 10} Further, the mesenchymal stem cell treatment that prevented the post-stroke induction of MMP-12 attenuated the post-stroke brain damage.^{24 25}

Post-stroke induction of MMP-8 (~ 279 -fold vs sham) was the largest after MMP-12 in mice. Our results are in agreement with a recent report wherein the authors




Figure 3 Alteration in the expression of matrix metalloproteinases (MMPs) under normal conditions in mice with genetic deletion of MMP-12. Real-time PCR analysis of the brain samples of healthy, young adult wild-type and MMP-12 knockout mice. Histograms and error bars indicate the mean and SD, respectively.

have demonstrated the upregulation of MMP-8 mRNA and protein in mouse ischaemic brains at 1-day reperfusion subsequent to a transient focal cerebral ischaemia.¹² Elevated MMP-8 brain mRNA expression was also recently reported at 3 and 7 days post-ischaemia induction in a photothrombotic ischaemic mouse model.¹⁵ Although the post-stroke induction of MMP-8 at 1 day after reperfusion in rats is minimal (~3-fold vs sham), the increase in mRNA expression is prominent (~35-fold vs sham at 5 and 7 days of reperfusion) at later reperfusion time points.¹⁰ Overall, the post-stroke induction of MMP-8 was evident in both the rodent stroke models. Post-stroke induction of MMP-8 was also reported in human brain.⁸ Inactivation of MMP-8 either by administration of MMP-8 inhibitor or MMP-8 specific shRNA immediately after reperfusion subsequent to a transient focal cerebral ischaemia reduced the extent of brain damage.¹²

At 1-day reperfusion after a transient focal cerebral ischaemia, MMP-3 was downregulated (~5-fold vs sham) in the ischaemic brains of mice. In contrast, in rats, mRNA expression was upregulated (~11-fold vs sham) along with protein expression at 1-day reperfusion; however no change was noticed in mRNA expression at 3, 5, and 7 days of reperfusion.^{6, 10} In addition, increased protein expression of MMP-3 was reported in reactive astrocytes and/or microglia/macrophages at 3 days of reperfusion in mice.²⁶ Increased MMP-3 protein expression was recently reported in brain tissue of deceased patients who had an ischaemic stroke within the previous 4 days.⁸ MMP-3 can proteolytically activate MMP-9.^{27, 28} The post-stroke induction of MMP-3 could be detrimental because its inhibition either by administration of its inhibitor or MMP-3 specific shRNA in rats significantly improved the functional outcome in hyperglycaemic stroke.¹¹

Upregulation of MMP-2 was reported in the ischaemic brain of rats at 5 days of reperfusion as compared with its expression at the initial time points after reperfusion.²⁹

In agreement with these findings, we reported a marked increase in MMP-2 activity in rats at 7 days of reperfusion.¹⁰ However, we did not notice a prominent difference in the mRNA expression of MMP-2 until 7 days of reperfusion. In contrast, in this study, MMP-2 was reasonably upregulated (~11-fold vs sham) in mice at 1-day reperfusion. Elevated MMP-2 brain mRNA expression was also recently reported at 3 and 7 days post-ischaemia induction in a photothrombotic ischaemic mouse model.¹⁵

Interestingly, MMP-1a, MMP-10, MMP-15, MMP-16, MMP-17 and MMP-21 were downregulated (>10-fold vs sham) in mice in this study in contrast to rats at 1-day reperfusion.¹⁰ Of all the MMPs that were downregulated in mice, the degree of reduction in MMP-10 mRNA expression (~247-fold vs sham) is predominant over other MMPs. In a thromboembolic stroke model, MMP-10 knockout mice had similar infarct size as the wild-type mice.⁹ Administration of MMP-10 to wild-type mice reduced the infarct size. In the same thromboembolic stroke model, administration of MMP-10 in combination with tissue-type plasminogen activator (tPA) significantly reduced the infarct size as compared with tPA alone.¹⁴ In contrast to animal studies, increased MMP-10 protein levels were reported in human ischaemic brain.⁸

Most of the studies we discussed here in mice with genetic deletion of certain MMPs are in conventional knockout models (constitutive). The constitutive knockout of a target MMP is likely to result in the compensatory changes in these mice with altered expression of other MMPs. This further complicates the interpretation of the data that will be generated from these animal models. In this study, genetic deletion of MMP-12 prevented the post-stroke upregulation of MMP-12, MMP-2 and MMP-8 and downregulation of MMP-1a, MMP-10 and MMP-21, as well as upregulated MMP-28 as compared with wild-type mice. In fact, the genetic deletion of MMP-12 under normal conditions also resulted in the altered expression

of several other MMPs. Our findings demonstrated that the MMP-12 knockout mice are not the suitable models to conclusively assess the role of MMP-12. It also raises the question of whether the same apply to other MMP deleted constitutive knockout mice.

In summary, our results demonstrate the differences in the post-stroke expression of MMPs in the ischaemic brain. Of all the MMPs, MMP-12 upregulation is predominant in the ischaemic brain of both rodent species. Because of the compensatory changes, constitutive MMP knockout mice may not be the suitable models for confirmatory studies.

Acknowledgements We thank Mary Allyson Finger for assistance in manuscript format and review.

Contributors KKV was involved in the conception and hypotheses delineation, design and execution of the study, analysis and interpretation of results. KRN was involved in animal handling, postsurgical animal care, collection of tissues, conduct of PCR analysis, acquisition and quantification of the data, statistical analysis and manuscript preparation. BC induced ischaemia in animals. IBM and DNF assisted KRN in PCR analysis. AM and IV assisted KRN in agarose gel electrophoresis. JD, DMP, KMB and KKV reviewed and edited the manuscript.

Funding This work was supported by the Illinois Neurological Institute, OSF HealthCare Foundation.

Competing interests None declared.

Patient consent Not required.

Ethics approval The Institutional Animal Care and Use Committee of the University of Illinois College of Medicine at Peoria.

Provenance and peer review Commissioned by the CSA; externally peer reviewed.

Data sharing statement The authors agree to provide upon request the unpublished data that include raw data, metadata and statistical analysis results for the purposes of reproducing the results.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Lee HS, Namkoong K, Kim DH, *et al.* Hydrogen peroxide-induced alterations of tight junction proteins in bovine brain microvascular endothelial cells. *Microvasc Res* 2004;68:231–8.
- Si-Tayeb K, Monvoisin A, Mazzocco C, *et al.* Matrix metalloproteinase 3 is present in the cell nucleus and is involved in apoptosis. *Am J Pathol* 2006;169:1390–401.
- Rosenberg GA, Yang Y. Vasogenic edema due to tight junction disruption by matrix metalloproteinases in cerebral ischemia. *Neurosurg Focus* 2007;22:1–9.
- Wasserman JK, Schlichter LC. Minocycline protects the blood–brain barrier and reduces edema following intracerebral hemorrhage in the rat. *Exp Neurol* 2007;207:227–37.
- Yang Y, Candelario-Jalil E, Thompson JF, *et al.* Increased intranuclear matrix metalloproteinase activity in neurons interferes with oxidative DNA repair in focal cerebral ischemia. *J Neurochem* 2010;112:134–49.
- Chelluboina B, Klopfenstein JD, Pinson DM, *et al.* Matrix metalloproteinase-12 induces blood–brain barrier damage after focal cerebral ischemia. *Stroke* 2015;46:3523–31.
- Zhao BQ, Wang S, Kim HY, *et al.* Role of matrix metalloproteinases in delayed cortical responses after stroke. *Nat Med* 2006;12:441–5.
- Cuadrado E, Rosell A, Penalba A, *et al.* Vascular MMP-9/TIMP-2 and neuronal MMP-10 up-regulation in human brain after stroke: a combined laser microdissection and protein array study. *J Proteome Res* 2009;8:3191–7.
- Orbe J, Barrenetxe J, Rodriguez JA, *et al.* Matrix metalloproteinase-10 effectively reduces infarct size in experimental stroke by enhancing fibrinolysis via a thrombin-activatable fibrinolysis inhibitor-mediated mechanism. *Circulation* 2011;124:2909–19.
- Chelluboina B, Warhekar A, Dillard M, *et al.* Post-transcriptional inactivation of matrix metalloproteinase-12 after focal cerebral ischemia attenuates brain damage. *Sci Rep* 2015;5:9504.
- Hafez S, Abdelsaid M, El-Shafey S, *et al.* Matrix metalloproteinase 3 exacerbates hemorrhagic transformation and worsens functional outcomes in hyperglycemic stroke. *Stroke* 2016;47:843–51.
- Han JE, Lee EJ, Moon E, *et al.* Matrix Metalloproteinase-8 is a Novel Pathogenic Factor in Focal Cerebral Ischemia. *Mol Neurobiol* 2016;53:231–9.
- Ma F, Martínez-San Segundo P, Barceló V, *et al.* Matrix metalloproteinase-13 participates in neuroprotection and neurorepair after cerebral ischemia in mice. *Neurobiol Dis* 2016;91:236–46.
- Roncal C, Martínez de Lizarrondo S, Salicio A, *et al.* New thrombolytic strategy providing neuroprotection in experimental ischemic stroke: MMP10 alone or in combination with tissue-type plasminogen activator. *Cardiovasc Res* 2017;113:1219–29.
- Hirono J, Sanaki H, Kitada K, *et al.* Expression of tissue inhibitor of metalloproteinases and matrix metalloproteinases in the ischemic brain of photothrombosis model mice. *Neuroreport* 2018;29:174–180.
- Chaturvedi M, Kaczmarek L. Mmp-9 inhibition: a therapeutic strategy in ischemic stroke. *Mol Neurobiol* 2014;49:563–73.
- Heo JH, Lucero J, Abumiya T, *et al.* Matrix metalloproteinases increase very early during experimental focal cerebral ischemia. *J Cereb Blood Flow Metab* 1999;19:624–33.
- Clark AW, Krekoski CA, Bou SS, *et al.* Increased gelatinase A (MMP-2) and gelatinase B (MMP-9) activities in human brain after focal ischemia. *Neurosci Lett* 1997;238:53–6.
- Rosell A, Ortega-Aznar A, Alvarez-Sabin J, *et al.* Increased brain expression of matrix metalloproteinase-9 after ischemic and hemorrhagic human stroke. *Stroke* 2006;37:1399–406.
- Wang X, Jung J, Asahi M, *et al.* Effects of matrix metalloproteinase-9 gene knock-out on morphological and motor outcomes after traumatic brain injury. *J Neurosci* 2000;20:7037–42.
- Asahi M, Wang X, Mori T, *et al.* Effects of matrix metalloproteinase-9 gene knock-out on the proteolysis of blood–brain barrier and white matter components after cerebral ischemia. *J Neurosci* 2001;21:7724–32.
- Svedin P, Hagberg H, Sävman K, *et al.* Matrix metalloproteinase-9 gene knock-out protects the immature brain after cerebral hypoxia–ischemia. *J Neurosci* 2007;27:1511–8.
- Matsumoto S, Kobayashi T, Katoh M, *et al.* Expression and localization of matrix metalloproteinase-12 in the aorta of cholesterol-fed rabbits: relationship to lesion development. *Am J Pathol* 1998;153:109–19.
- Chelluboina B, Klopfenstein JD, Pinson DM, *et al.* Stem cell treatment after cerebral ischemia regulates the gene expression of apoptotic molecules. *Neurochem Res* 2014;39:1511–21.
- Chelluboina B, Nalamolu KR, Mendez GG, *et al.* Mesenchymal stem cell treatment prevents post-stroke dysregulation of matrix metalloproteinases and tissue inhibitors of metalloproteinases. *Cell Physiol Biochem* 2017;44:1360–9.
- Yang Y, Jalal FY, Thompson JF, *et al.* Tissue inhibitor of metalloproteinases-3 mediates the death of immature oligodendrocytes via TNF- α /TACE in focal cerebral ischemia in mice. *J Neuroinflammation* 2011;8:108.
- Candelario-Jalil E, Yang Y, Rosenberg GA. Diverse roles of matrix metalloproteinases and tissue inhibitors of metalloproteinases in neuroinflammation and cerebral ischemia. *Neuroscience* 2009;158:983–94.
- Yang Y, Hill JW, Rosenberg GA. Multiple roles of metalloproteinases in neurological disorders. *Prog Mol Biol Transl Sci* 2011;99:241–63.
- Rosenberg GA, Estrada EY, Dencoff JE. Matrix metalloproteinases and TIMPs are associated with blood–brain barrier opening after reperfusion in rat brain. *Stroke* 1998;29:2189–95.