

SUPPLEMENTAL TABLES**Table 1.** Baseline characteristics of patients included versus not included in this substudy of CHANCE trial.

Characteristics	Patients included	All other patients	P Value
	(n=807)	(n=4363)	
Age, y, median (IQR)	63.1(55.0-72.0)	62.1(54.7-71.2)	0.10
Male, n (%)	525(65.1)	2,895(66.4)	0.47
Medical history, n (%)			
Ischaemic stroke	140(17.3)	893(20.5)	0.04
TIA	26(3.2)	148(3.4)	0.81
Myocardial infarction	14(1.7)	82(1.9)	0.78
Known atrial fibrillation or flutter	18(2.2)	78(1.8)	0.39
Angina	23(2.9)	161(3.7)	0.24
Valvular heart disease	3(0.4)	11(0.3)	0.55
Hypertension	527(65.3)	2872(65.8)	0.77
Diabetes mellitus	155(19.2)	938(21.5)	0.14
Hypercholesterolaemia	105(13.0)	468(10.7)	0.06
Current or previous smoking, n (%)	343(42.5)	1878(43.0)	0.78
Time to randomization, n (%)			0.14
<12 hours	421(52.2)	2152(49.3)	
≥ 12hours	386(47.8)	2211(50.7)	
NIHSS on admission, median (IQR)	2(0-2)	1(0-2)	0.02
Qualifying event, n (%)			0.04
TIA	202(25.0)	1243(28.5)	
Minor stroke	605(75.0)	3120(71.5)	
Group, n (%)			0.98
Aspirin only	404(50.1)	2182(50.0)	

Clopidogrel+aspirin	403(49.9)	2181(50.0)	
Medications, n (%)			
Antihypertensive	359(44.7)	1455(33.6)	<0.001
Antidiabetic	110(13.7)	546(12.6)	0.40
Lipid-lowering	423(52.6)	1748(40.4)	<0.001
Outcomes			
Ischaemic stroke	84(10.4)	528(12.1)	0.17
Composite events ^a	89(11.0)	563(12.9)	0.14
TIA	25(3.1)	94(2.2)	0.10
Any bleeding	18(2.2)	92(2.1)	0.83

Abbreviations: TIA: transient ischaemic attack; NAI: no acute infarction; SAI: single acute infarction; MAIs: multiple acute infarctions; IQR: interquartile range.

^a Composite events: stroke, myocardial infarction, or death from cardiovascular causes.