

CO₂ combining power and outcomes in patients with acute ischaemic stroke or transient ischaemic attack

Anxin Wang ^{1,2}, Xue Tian,^{3,4} Hongqiu Gu ^{1,2}, Yingting Zuo,^{3,4} Xia Meng,^{1,2} Wei Lv,^{1,2} Hao Li ^{1,2}, Yongjun Wang ^{1,2}

To cite: Wang A, Tian X, Gu H, *et al.* CO₂ combining power and outcomes in patients with acute ischaemic stroke or transient ischaemic attack. *Stroke & Vascular Neurology* 2020;0. doi:10.1136/svn-2020-000476

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/svn-2020-000476>).

Received 1 July 2020

Revised 13 October 2020

Accepted 14 November 2020



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹China National Clinical Research Center for Neurological Diseases, Beijing Tiantan Hospital, Capital Medical University, Beijing, China

²Department of Neurology, Beijing Tiantan Hospital, Capital Medical University, Beijing, China

³Department of Epidemiology and Health Statistics, School of Public Health, Capital Medical University, Beijing, China

⁴Beijing Municipal Key Laboratory of Clinical Epidemiology, Beijing, China

Correspondence to

Dr Yongjun Wang, Department of Neurology, ; yongjunwang@nccrncd.org.cn

ABSTRACT

Background and purpose The clinical significance of carbon dioxide combining power (CO₂CP) in ischaemic cerebrovascular disease is not well established, and the role of CO₂CP in the prognosis of acute ischaemic stroke (AIS) or transient ischaemic attack (TIA) has not been reported. The objective of the study was to investigate the associations between CO₂CP and clinical outcomes in patients with AIS or TIA.

Methods Data were derived from the China National Stroke Registry III. Patients were classified into five groups by quintiles of CO₂CP levels and three groups according to the normal range of CO₂CP (23–29 mmol/L). Multivariable Cox and logistic regressions were adopted to explore the associations of CO₂CP levels with all-cause death and poor functional outcomes (modified Rankin Scale (mRS) 3–6/2–6) at 3 months and 1 year.

Results Among 9531 patients included in the study, the median (IQR) CO₂CP was 24.9 (23.0–27.0) mmol/L. After adjustment for potential confounders, patients in the first CO₂CP quintile (21.1–23.3 mmol/L) had higher risk of all-cause death and poor functional outcomes (mRS score of 3–6/2–6) (HR or OR with 95% CI 2.37 (1.32 to 4.25), 1.49 (1.20 to 1.83) and 1.21 (1.03 to 1.42), respectively) compared with those in the fourth quintile. Similar results were found for outcomes at 1 year. Furthermore, all associations were also significant when CO₂CP was <23 mmol/L compared with CO₂CP of 23–29 mmol/L.

Conclusions Decreased CO₂CP was associated with high risk of all-cause death and poor functional outcomes in patients with AIS or TIA.

INTRODUCTION

Carbon dioxide combining power (CO₂CP) is a measure of alkali reserve and can help to diagnose the metabolic type of acidosis and alkalosis.^{1,2} The determination of CO₂CP basically represents the amount of alkali reserves in the blood and determines whether an acid-base disturbance is present and if so to what degree. Decreased CO₂CP has been demonstrated to be an indicator of metabolic acidosis in patients without respiratory disease.²

Previous studies have shown that metabolic acidosis is associated with insulin resistance,³ hypertension,⁴ inflammation,⁵ activation of the renin-angiotensin-aldosterone system⁶ and endothelial dysfunction,⁷ all of which can

contribute to adverse clinical outcomes after stroke.^{3,8} Furthermore, acidosis is common and has been proven to be associated with poor outcomes in ischaemic stroke.⁹ Herein, it is logical to consider CO₂CP as a risk factor for poor outcomes of stroke.

Clinical evidence has shown that decreased CO₂CP is associated with advanced clinical stages of cancer and may predict worse outcomes of disease-free survival in patients with stage II/III colorectal cancer.¹ Another study revealed that decreased CO₂CP was an independent risk factor for the development of acute kidney injury and in-hospital mortality. However, the role of CO₂CP in the prognosis of acute ischaemic stroke (AIS) or transient ischaemic attack (TIA) has not been examined to date. Therefore, in the present study, we sought to investigate the potential associations between CO₂CP levels and adverse clinical outcomes in patients with AIS or TIA.

METHODS

Study population

The data analysed in this study were obtained from the Third China National Stroke Registry (CNSR-III), which is a nationwide prospective registry including 201 hospitals of 22 provinces. The registry contains data of patients with AIS or TIA who presented to these hospitals from August 2015 to March 2018. A total of 15 166 participants with AIS or TIA at 7 days from symptom onset were enrolled in CNSR-III. The details, rationale and basic description of CNSR-III have been published previously.¹⁰

Data collection

Baseline data were prospectively collected using an electronic data capture system by face-to-face interviews following a standard data collection protocol that was developed by the steering committee, which included age, sex, body mass index (BMI; calculated as

weight in kilograms divided by height in metres squared, kg/m²), medical history (hypertension, diabetes mellitus, dyslipidaemia, stroke or TIA, atrial fibrillation or flutter, peripheral vascular disease, heart failure, chronic obstructive pulmonary disease (COPD)), stroke type (ischemic stroke or TIA), aetiology classification of ischaemic stroke performed according to the TOAST (Trial of Org 10172 in Acute Stroke Treatment) criteria,¹¹ current smoker, in-hospital medication (cholesterol-lowering agents, anti-hypertensive agents, hypoglycaemic agents, antiplatelet agents and anticoagulant agents), severity of stroke on admission (National Institutes of Health Stroke Scale, NIHSS),¹² time from onset of symptoms to admission, total cholesterol, low-density lipoprotein cholesterol, high-density lipoprotein cholesterol, triglyceride, fasting blood glucose, estimated glomerular filtration rate calculated using the creatinine-based Chronic Kidney Disease Epidemiological Collaboration equation¹³ and high-sensitivity C reactive protein (hs-CRP), and medication on discharge (cholesterol-lowering agents, antihypertensive agents, hypoglycaemic agents, antiplatelet agents and anticoagulant agents).

CO₂CP testing

Fasting blood samples were collected at 24 hours after admission. The samples were placed into serum-separation tubes and EDTA anticoagulation blood collection tube and kept at room temperature. Afterwards, CO₂CP was analysed by automated haematology analyser at 1 hour after sample collection at each research centre. All measurements were performed by laboratory personnel blinded to subjects' clinical situations. CO₂CP was analysed as a continuous variable and divided into quintiles.

Outcome assessment

Patients were followed up for clinical outcomes at 3 months and 1 year after symptom onset. They were interviewed face to face at 3 months and contacted over the telephone by trained research coordinators at 1 year. Data regarding all-cause death and poor functional outcome were collected by trained research coordinators who were blinded to subjects' baseline characteristics. All-cause death was defined as death from any cause and confirmed by a death certification from the attended hospital or the local citizen registry. Poor functional outcome was defined as modified Rankin Scale (mRS) score ranging from 2 to 6 or from 3 to 6 at 3 months and 1 year.

Statistical analysis

Patients were classified into five groups by CO₂CP quintiles. Continuous variables are described as median and IQR due to skewed distribution, and categorical variables are described as frequencies and percentages. Non-parametric Wilcoxon or Kruskal-Wallis test was used to compare group differences for continuous variables, and χ^2 test or Fisher's exact test was used for categorical variables.

We took the fourth quintile with the lowest incidence of outcomes as a reference and performed Cox proportional hazard models and logistic regression to explore the associations of CO₂CP with all-cause death and poor functional outcomes, respectively. Because 201 hospitals participated in the study, the hospitals were treated as clusters in the model and the sandwich estimated was used to account for the correlations. Variables were adjusted in the multivariable analyses if associated with CO₂CP in the univariate analysis with $p < 0.2$. Adjusted HRs or ORs and their 95% CIs were calculated. We used three adjusted models. Model 1 was adjusted for age and gender; model 2 was further adjusted for BMI, medical history (IS, TIA, hyperlipidaemia and atrial fibrillation or flutter), stroke type, current smoker, antihypertensive agents, anticoagulant agents, NIHSS score and high-density lipoprotein cholesterol on admission; and model 3 was further adjusted for history of COPD, time from onset of symptoms to admission and hs-CRP on admission. Trend tests were performed in the regression models after the median CO₂CP values of each quartile were entered into the model and treated as a continuous variable. Multivariate survival analysis was performed by the Kaplan-Meier method. In addition, we used restricted cubic splines to examine the shape of the association between CO₂CP and outcomes with five knots (at the 5th, 25th, 50th, 75th and 95th percentiles); the reference point for CO₂CP was the median of the reference group (the fourth quintile), and the HR/OR was adjusted for all confounding variables. Subgroup analyses were performed stratified by age (<65 and ≥ 65 years), gender, stroke subtype (IS and TIA), smoking status (yes and no) and NIHSS score (<4 and ≥ 4).

To further investigate the associations of CO₂CP levels and outcomes, patients were classified into three groups (<23, 23–29 and ≥ 29 mmol/L) according to the normal range of CO₂CP levels (23–29 mmol/L).² Additionally, we also used C statistics, integrated discrimination (IDI) and net reclassification index (NRI) to evaluate the incremental predictive value of CO₂CP beyond conventional risk factors.

All analyses were performed with SAS V.9.4 software. A two-tailed value of $p < 0.05$ was considered statistically significant.

RESULTS

Baseline characteristics

After exclusion of patients with missing data on CO₂CP ($n=5339$) and mRS at 3 months or 1 year ($n=296$), 9531 patients were included in our analysis. Baseline comparison of the excluded and included patients is presented in online supplemental table S1.

The baseline characteristics of included patients stratified into quintiles according to CO₂CP are shown in [table 1](#). The median (IQR) age of the patients was 62 (54–70) years, 6556 (68.79%) were men, and the median (IQR) CO₂CP level was 24.9 (23.0–27.0) mmol/L.

Table 1 Baseline characteristics of included patients stratified into quintiles according to CO₂CP

Variable	Quintiles of CO ₂ CP, mmol/L					P value	
	Overall	Q1	Q2	Q3	Q4		Q5
n	9531	1914 (20.08)	1956 (20.52)	1874 (19.66)	1869 (19.61)	1918 (20.12)	
CO ₂ CP, mmol/L	24.9 (23.0–27.0)	21.1 (20.0–22.0)	23.3 (23.0–23.8)	25.0 (24.5–25.2)	26.4 (26.0–27.0)	28.8 (28.0–30.0)	<0.0001
Age, years	62 (54–70)	62 (53–70)	62 (54–70)	62 (54–69)	62 (55–70)	63 (56–70)	0.0008
Men, n (%)	6556 (68.79)	1347 (70.38)	1308 (66.87)	1319 (70.38)	1278 (68.38)	1304 (67.99)	0.0738
Body mass index, kg/m ²	24.51 (22.68–26.67)	24.66 (22.86–26.71)	24.62 (22.86–26.7)	24.49 (22.68–26.67)	24.57 (22.66–26.57)	24.49 (22.49–26.64)	0.1261
Medical history, n (%)							
Hypertension	5958 (62.51)	1170 (61.13)	1249 (63.85)	1190 (63.50)	1149 (61.48)	1200 (62.57)	0.3187
Diabetes mellitus	2243 (23.53)	421 (22.00)	456 (23.31)	452 (24.12)	459 (24.56)	455 (23.72)	0.3986
Dyslipidaemia	801 (8.4)	133 (6.95)	156 (7.98)	160 (8.54)	178 (9.52)	174 (9.07)	0.0416
Stroke or TIA	2190 (22.98)	393 (20.53)	447 (22.85)	436 (23.27)	434 (23.22)	480 (25.03)	0.0246
Atrial fibrillation/flutter	621 (6.52)	152 (7.94)	123 (6.29)	125 (6.67)	109 (5.83)	112 (5.84)	0.0497
Peripheral vascular disease	66 (0.69)	14 (0.73)	9 (0.46)	13 (0.69)	16 (0.86)	14 (0.73)	0.6727
Heart failure	64 (4.81)	14 (5.51)	15 (5.66)	12 (4.44)	9 (3.46)	14 (4.96)	0.7691
COPD	81 (0.85)	15 (0.78)	13 (0.66)	16 (0.85)	12 (0.64)	25 (1.30)	0.1625
Stroke type/subtype, n (%)							
Ischaemic stroke	8832 (92.67)	1789 (93.47)	1820 (93.05)	1735 (92.58)	1739 (93.04)	1749 (91.19)	0.0661
Transient ischaemic attack	699 (7.33)	125 (6.53)	136 (6.95)	139 (7.42)	130 (6.96)	169 (8.81)	
TOAST, n (%)							
Large-artery atherosclerosis	2400 (25.18)	520 (27.17)	477 (24.39)	486 (25.93)	452 (24.18)	465 (24.24)	0.3994
Cardioembolism	544 (5.71)	123 (6.43)	112 (5.73)	104 (5.55)	99 (5.30)	106 (5.53)	
Small-vessel occlusion	1966 (20.63)	355 (18.55)	396 (20.25)	391 (20.86)	402 (21.51)	422 (22.00)	
Other determined aetiology	126 (1.32)	26 (1.36)	30 (1.53)	22 (1.17)	26 (1.39)	22 (1.15)	
Undetermined aetiology	4495 (47.16)	890 (46.50)	941 (48.11)	871 (46.48)	890 (47.62)	903 (47.08)	
Current smoker, n (%)	3089 (32.41)	684 (35.74)	634 (32.41)	621 (33.14)	577 (30.87)	573 (29.87)	0.0099
Medication in hospital, n (%)							
Cholesterol-lowering agents	9182 (96.99)	1844 (97.10)	1893 (97.23)	1803 (96.99)	1791 (96.45)	1851 (97.17)	0.6374
Antihypertensive agents	4568 (48.25)	959 (50.50)	939 (48.23)	874 (47.01)	873 (47.01)	923 (48.45)	0.1875
Hypoglycaemic agents	2449 (25.87)	476 (25.07)	506 (25.99)	489 (26.30)	486 (26.17)	492 (25.83)	0.9206
Antiplatelet agents	9191 (97.08)	1852 (97.53)	1885 (96.82)	1806 (97.15)	1811 (97.52)	1837 (96.43)	0.2016
Anticoagulant agents	849 (8.97)	211 (11.11)	177 (9.09)	159 (8.55)	150 (8.08)	152 (7.98)	0.0043
NIHSS score on admission	3 (1–6)	3 (1–6)	3 (1–6)	3 (1–6)	3 (1–5)	3 (1–6)	<0.0001
Time from onset of symptoms to admission, hours	15.00 (3.00–45.00)	12.00 (3.00–32.00)	14.00 (3.00–45.00)	15.00 (3.00–45.00)	15.00 (3.00–44.00)	19.00 (4.00–48.00)	<0.0001

Continued

Table 1 Continued

Variable	Overall	Quintiles of CO ₂ CP, mmol/L					P value
		Q1	Q2	Q3	Q4	Q5	
Total cholesterol, mmol/L	3.97 (3.31–4.74)	3.92 (3.31–4.74)	4.02 (3.31–4.77)	3.95 (3.3–4.71)	4 (3.31–4.74)	3.95 (3.32–4.75)	0.7846
Low-density lipoprotein, mmol/L	2.3 (1.71–2.97)	2.27 (1.7–2.96)	2.34 (1.72–3.01)	2.28 (1.69–2.95)	2.31 (1.71–2.97)	2.3 (1.72–2.98)	0.6915
High-density lipoprotein, mmol/L	0.93 (0.77–1.11)	0.90 (0.77–1.08)	0.93 (0.76–1.11)	0.93 (0.76–1.1)	0.93 (0.77–1.13)	0.95 (0.78–1.13)	0.0236
Triglycerides, mmol/L	1.38 (1.04–1.89)	1.38 (1.04–1.91)	1.40 (1.06–1.91)	1.36 (1.00–1.88)	1.41 (1.04–1.95)	1.36 (1.04–1.84)	0.2033
Fasting blood glucose, mmol/L	5.56 (4.90–6.99)	5.53 (4.87–6.94)	5.60 (4.94–7.10)	5.56 (4.90–6.90)	5.52 (4.90–7.02)	5.53 (4.90–6.99)	0.3506
eGFR, mL/min/1.73 m ²	93.31 (81.94–102.03)	92.96 (80.93–102.93)	93.32 (81.58–101.83)	93.85 (81.95–102.55)	94.11 (82.57–102.01)	92.76 (82.72–101.01)	0.4400
hs-CRP, mg/L	1.78 (0.81–4.71)	2.05 (0.87–5.58)	1.79 (0.82–4.78)	1.75 (0.8–4.51)	1.66 (0.81–4.43)	1.64 (0.78–4.36)	0.0020
Medication on discharge, n (%)							
Cholesterol-lowering agents	8571 (92.06)	1772 (93.07)	1801 (92.26)	1717 (91.82)	1709 (91.59)	1752 (91.49)	0.3638
Antihypertensive agents	4840 (50.91)	1004 (52.73)	996 (51.02)	934 (49.95)	944 (50.59)	962 (50.23)	0.4544
Hypoglycaemic agents	2298 (24.17)	460 (24.16)	478 (24.49)	449 (24.01)	445 (23.85)	466 (24.33)	0.9918
Antiplatelet agents	8668 (91.17)	1740 (91.93)	1773 (90.83)	1710 (91.44)	1704 (91.32)	1741 (90.91)	0.9421
Anticoagulant agents	260 (2.73)	64 (3.36)	53 (2.72)	47 (2.51)	44 (2.36)	52 (2.72)	0.3853

CO₂CP, carbon dioxide combining power; COPD, chronic obstructive pulmonary disease; eGFR, estimated glomerular filtration rate; hs-CRP, high-sensitivity C reactive protein; NIHSS, National Institutes of Health Stroke Scale; TIA, transient ischaemic attack; TOAST, Trial of Org 10172 in Acute Stroke Treatment.

Table 2 Association of all-cause death and poor functional outcomes with quintiles of CO₂CP

Outcomes	Quintile of CO ₂ CP					P for trend
	Q1	Q2	Q3	Q4	Q5	
At 3 months						
Death	39 (2.04)	27 (1.38)	21 (1.12)	16 (0.86)	23 (1.20)	
Unadjusted	2.55 (1.37–4.78)	1.77 (0.94–3.33)	1.40 (0.73–2.72)	Reference	1.24 (0.64–2.39)	0.7585
Model 1	2.65 (1.41–4.96)	1.70 (0.89–3.22)	1.41 (0.73–2.75)	Reference	1.16 (0.60–2.27)	0.6639
Model 2	2.33 (1.23–4.44)	1.63 (0.85–3.13)	1.24 (0.62–2.46)	Reference	1.15 (0.58–2.27)	0.7527
Model 3	2.40 (1.27–4.55)	1.58 (0.82–3.03)	1.16 (0.58–2.32)	Reference	1.29 (0.64–2.57)	0.8246
mRS 3–6	319 (16.67)	276 (14.11)	257 (13.71)	208 (11.13)	228 (11.89)	
Unadjusted	1.74 (1.41–2.14)	1.39 (1.13–1.70)	1.30 (1.06–1.59)	Reference	1.05 (0.85–1.29)	0.5259
Model 1	1.80 (1.46–2.22)	1.40 (1.14–1.72)	1.34 (1.09–1.65)	Reference	1.02 (0.83–1.26)	0.3626
Model 2	1.57 (1.25–1.97)	1.27 (1.01–1.58)	1.25 (1.00–1.57)	Reference	0.96 (0.76–1.21)	0.3243
Model 3	1.58 (1.26–1.98)	1.27 (1.02–1.59)	1.27 (1.01–1.59)	Reference	0.97 (0.77–1.23)	0.3814
mRS 2–6	561 (29.31)	516 (26.38)	483 (25.77)	448 (23.97)	476 (24.82)	
Unadjusted	1.41 (1.20–1.66)	1.18 (1.01–1.38)	1.11 (0.95–1.30)	Reference	1.07 (0.91–1.25)	0.5389
Model 1	1.46 (1.24–1.72)	1.19 (1.02–1.40)	1.14 (0.97–1.34)	Reference	1.05 (0.90–1.24)	0.3668
Model 2	1.26 (1.05–1.51)	1.09 (0.92–1.30)	1.07 (0.90–1.28)	Reference	1.02 (0.85–1.21)	0.4684
Model 3	1.26 (1.05–1.51)	1.10 (0.92–1.30)	1.08 (0.91–1.28)	Reference	1.02 (0.86–1.22)	0.4954
At 1 year						
Death	70 (3.66)	60 (3.07)	58 (3.09)	41 (2.19)	51 (2.66)	
Unadjusted	1.64 (1.08–2.48)	1.41 (0.94–2.12)	1.42 (0.95–2.14)	Reference	1.11 (0.72–1.70)	0.7947
Model 1	1.68 (1.11–2.54)	1.40 (0.93–2.11)	1.48 (0.98–2.22)	Reference	1.05 (0.68–1.62)	0.9283
Model 2	1.50 (0.99–2.28)	1.35 (0.89–2.04)	1.34 (0.89–2.03)	Reference	1.00 (0.65–1.55)	0.9980
Model 3	1.56 (1.03–2.36)	1.35 (0.89–2.04)	1.35 (0.89–2.04)	Reference	1.06 (0.68–1.65)	0.8666
mRS 3–6	301 (15.73)	243 (12.42)	246 (13.13)	212 (11.34)	233 (12.15)	
Unadjusted	1.44 (1.17–1.77)	1.11 (0.91–1.37)	1.16 (0.95–1.42)	Reference	1.05 (0.86–1.30)	0.6571
Model 1	1.50 (1.21–1.85)	1.12 (0.91–1.38)	1.21 (0.98–1.49)	Reference	1.03 (0.83–1.27)	0.4322
Model 2	1.32 (1.05–1.66)	1.01 (0.81–1.27)	1.11 (0.88–1.38)	Reference	0.97 (0.77–1.22)	0.3000
Model 3	1.33 (1.06–1.67)	1.01 (0.81–1.27)	1.12 (0.89–1.41)	Reference	0.98 (0.78–1.24)	0.3659
mRS 2–6	520 (27.17)	472 (24.13)	455 (24.28)	401 (21.46)	441 (22.99)	
Unadjusted	1.47 (1.24–1.73)	1.21 (1.03–1.43)	1.18 (1.01–1.38)	Reference	1.09 (0.93–1.29)	0.9855
Model 1	1.53 (1.30–1.81)	1.23 (1.05–1.45)	1.22 (1.04–1.44)	Reference	1.07 (0.91–1.27)	0.7330
Model 2	1.35 (1.13–1.62)	1.14 (0.96–1.36)	1.16 (0.97–1.38)	Reference	1.04 (0.87–1.25)	0.8040
Model 3	1.36 (1.13–1.62)	1.14 (0.96–1.36)	1.17 (0.98–1.39)	Reference	1.05 (0.88–1.25)	0.8458

Model 1: adjusted for age and gender.

Model 2: adjusted for age, gender, body mass index, history of dyslipidaemia, stroke or transient ischaemic stroke, atrial fibrillation/flutter, stroke type, current smoker, antihypertensive agents, anticoagulant agents, National Institutes of Health Stroke Scale score and high-density lipoprotein cholesterol on admission.

Model 3: adjusted for variables in model 2, plus history of chronic obstructive pulmonary disease, time from onset of symptoms to admission and high-sensitivity C reactive protein on admission.

CO₂CP, carbon dioxide combining power; mRS, modified Rankin Scale.

Compared with the rest of the quintiles, participants in the first quintile of CO₂CP (21.1–23.3 mmol/L) had higher proportions of atrial fibrillation or flutter, more current smokers, more anticoagulant agent takers, a shorter time from onset of symptoms to admission and higher hs-CRP levels. On the other hand, participants in the fifth quintile (≥28.8 mmol/L) were more likely to be older, have higher proportions of dyslipidaemia

and stroke, and have higher high-density lipoprotein cholesterol levels. In addition, patients in the fifth quintile were likely to have severe neurological deficits than those in the fourth quintile.

Associations of CO₂CP with all-cause death and poor functional outcome

At 3-month assessment, 126 (1.32%) patients had died, 1288 (13.51) patients had poor functional outcome




Figure 1 Kaplan-Meier curve of all-cause death incidence rate by quintiles of carbon dioxide combining power. (A) Death at 3 months and (B) death at 1 year. Adjusted for age, gender, body mass index, history of dyslipidaemia, stroke or transient ischaemic stroke, atrial fibrillation/flutter, stroke type, current smoker, antihypertensive agents, anticoagulant agents, National Institutes of Health Stroke Scale score, high-density lipoprotein cholesterol, history of chronic obstructive pulmonary disease, time from onset of symptoms to admission and high-sensitivity C reactive protein on admission.

(mRS score 3–6) and 2484 (26.06%) patients had mRS score 2–6. At 1-year follow-up, 280 (2.94%) patients had died, 1235 (12.96%) patients had mRS score 3–6 and 2289 (24.02%) patients had mRS score 2–6 (table 2). Kaplan-Meier curves by quintiles of CO₂CP showed that patients in the first quintile had a higher incidence of all-cause death both at 3 months and at 1 year ($p < 0.05$) (figure 1A,B).

After adjustment for the above-mentioned confounding variables, patients in the first quintile CO₂CP group were significantly associated with increased risk of all-cause death and poor functional outcome, compared with the fourth quintile of CO₂CP taken as reference; the adjusted HR/OR (95% CI) was 2.37 (1.32 to 4.25), 1.49 (1.20 to 1.83) and 1.21 (1.03 to 1.42) for all-cause death, mRS score 3–6 and mRS score 2–6 at 3 months, respectively. The associations remained significant at 1 year (table 2).

All the associations were consistent in prespecified subgroups (all p for interactions > 0.05 ; online supplemental table S2).

Multivariable-adjusted spline regression models showed that the relationships between CO₂CP and adverse outcomes were non-linear. A lower CO₂CP was associated with a higher risk of all-cause death at both 3-month and 1-year follow-up; as the CO₂CP level increased, the HR of all-cause death declined steadily, showing a somewhat L-shaped associations. Furthermore, similar-shaped associations also existed for mRS score 2–6 and mRS score 3–6 at 3 months and 1 year (figure 2A–F).

In the analysis of the three groups, patients with CO₂CP < 23 mmol/L had higher proportion of atrial fibrillation or flutter and anticoagulant agent takers and higher hs-CRP levels compared with other groups. On the other hand, those with CO₂CP ≥ 29 mmol/L were older and had higher proportion of stroke or TIA and less time from onset of symptoms to admission (online supplemental table S3). Furthermore, we found that patients with CO₂CP < 23 mmol/L had 64%, 26% and

16% higher risk of all-cause death and poor functional outcome (mRS score 3–6 and mRS score 2–6) compared with patients with normal CO₂CP (23–29 mmol/L) at 3-month follow-up; the adjusted HR/OR (95% CI) was 1.64 (1.08 to 2.49), 1.26 (1.08 to 1.48) and 1.16 (1.02 to 1.32), respectively. A similar trend was observed at 1-year follow-up (figure 3). However, there were no significant associations between CO₂CP and outcomes in patients with CO₂CP ≥ 29 mmol/L.

Incremental predictive value of CO₂CP

We evaluated whether CO₂CP would further increase the predictive value of conventional risk factors (online supplemental table S4). For death at 3 months as the outcome of interest, the C statistics by the conventional model did not significantly improve with the addition of CO₂CP (from 0.841 to 0.843). However, the discriminatory power and risk reclassification appeared to be substantially better (IDI 0.50%, $p = 0.0254$; continuous NRI = 19.43%, $p = 0.0303$). Similar results were found in poor functional outcomes and when the time point was set as 1 year.




Figure 2 Association of CO₂CP levels with risk of all-cause death and poor functional outcome at 3 months and 1 year. (A–C) All-cause death, mRS score 3–6 and mRS score 2–6 at 3 months; (D–F) all-cause death, mRS score 3–6 and mRS score 2–6 at 1 year. Adjusted for age, gender, body mass index, history of dyslipidaemia, stroke or transient ischaemic stroke, atrial fibrillation/flutter, stroke type, current smoker, antihypertensive agents, anticoagulant agents, National Institutes of Health Stroke Scale score, high-density lipoprotein cholesterol, history of chronic obstructive pulmonary disease, time from onset of symptoms to admission and high-sensitivity C reactive protein on admission. CO₂CP, carbon dioxide combining power; mRS, modified Rankin Scale.




Figure 3 Associations of CO₂CP levels stratified into three groups with the risk of all-cause death and poor functional outcome at 3 months and 1 year. Adjusted for age, gender, body mass index, history of dyslipidaemia, stroke or transient ischaemic stroke, atrial fibrillation/flutter, stroke type, current smoker, antihypertensive agents, anticoagulant agents, National Institutes of Health Stroke Scale score, high-density lipoprotein cholesterol, history of chronic obstructive pulmonary disease, time from onset of symptoms to admission and high-sensitivity C reactive protein on admission. CO₂CP, carbon dioxide combining power; mRS, modified Rankin Scale.

DISCUSSION

The major finding of the study was that decreased CO₂CP was significantly associated with risk of all-cause death and poor functional outcome in patients with AIS and TIA at 3 months and 1 year. Further analysis showed that patients with CO₂CP <23 mmol/L had high risk of adverse clinical outcomes, indicating that CO₂CP of 23 mmol/L may be an important point to discriminate all-cause death and poor functional outcomes in patients with AIS or TIA in clinical practice.

In accordance with previous findings that decreased CO₂CP is correlated with serious conditions and a poor disease prognosis,¹⁴⁻¹⁷ the findings of the present study showed that decreased CO₂CP was significantly associated with all-cause death and poor functional outcomes after stroke. The clinical implications of CO₂CP in the pathological conditions are mainly related to the acid-base disturbance, and studies have shown that the CO₂CP level can aid in the diagnosis of the metabolic types of acidosis and alkalosis. Decreased CO₂CP indicates insufficient alkali reserve and metabolic acidosis. Clinical data have shown that acidic pH is associated with an increased risk of mortality and unfavourable outcomes in patients with severe traumatic brain injury and ischaemic stroke.⁹

Moreover, an observational cohort study showed that metabolic acidosis was positively associated with a higher risk of ischaemic events and all-cause mortality in patients undergoing kidney transplantation.¹⁴ Results of several large cohort studies have further demonstrated that more severe metabolic acidosis is associated with higher mortality.¹⁵⁻¹⁷

Although most often decreased CO₂CP signifies the presence of metabolic acidosis, it may also reflect a decline in the bicarbonate concentration as compensation for respiratory alkalosis, which is a disturbance in the acid-base balance due to alveolar hyperventilation and is the most common acid-base disturbance observed in patients who are critically ill. Respiratory alkalosis is considered benign by many clinicians, but it can also be associated with a significant increase in poor prognosis and mortality.¹⁸ Experimental studies have revealed that a risk in brain pH induced by decreased CO₂CP impairs cortical GABAergic neurons, subsequently leading to vasoconstriction and deterioration of brain functions after transient ischaemia.^{19, 20} Furthermore, respiratory alkalosis was reported to precipitate cardiac arrhythmias and angina, reduced myocardial oxygen extraction and increased mortality according to population-based studies.^{18, 21, 22} All of these changes can lead to adverse clinical outcomes.

There are several plausible explanations for the association between CO₂CP and functional outcomes after AIS or TIA. First, metabolic acidosis induces insulin resistance, which can induce the production of proinflammatory cytokines in the brain after brain ischaemia.^{3, 23} Second, metabolic acidosis is also associated with inflammation, oxidative stress and malnutrition, which may increase the risk of mortality.^{24, 25} Third, acidosis is associated with toxic calcium influx into the cell and programmed cell death after ischaemic stroke.⁸ Because evidence regarding the associations between CO₂CP and clinical outcomes in patients with stroke is limited, further investigation is needed to clarify the underlying mechanism.

Strengths and limitations

The strengths of the study include the use of a multi-centre prospective registry with a large sample size, which resulted in sufficient statistical power. However, our study also had some limitations. First, this study only monitored baseline CO₂CP levels and did not examine the dynamic changes in CO₂CP, which may have provided more valuable information regarding the mechanisms underlying the associations. Second, heterogeneity of the equipment at different research centres may lead to biased estimates of results. However, this may have little impact because the hospitals were treated as clusters in the model and the sandwich estimated was used to account for the correlations in our study. Finally, some unmeasured or residual confounding effects may still exist due to the nature of the observational study.

CONCLUSIONS

Decreased CO₂CP was associated with high risk of all-cause death and poor functional outcome in patients with AIS or TIA at 3 months and 1 year. These associations were significant when CO₂CP level was <23 mmol/L. This finding underscores the importance of CO₂CP in the prognosis of AIS or TIA in clinical practice.

Acknowledgements We thank all participating hospitals, their physicians and nurses, CNSR III Steering Committee members and all the participants of the present study.

Contributors YW contributed to the conception and design of the study. AW contributed to manuscript drafting. HG, XT and YZ contributed to statistical analysis. XM and WL contributed to acquisition of data. HL contributed to critical revisions of the manuscript.

Funding This work was supported by the Young Elite Scientists Sponsorship Program by CAST (2018QNRC001), the Beijing Municipal Administration of Hospitals Incubating Program (PX2020021), the Beijing Municipal Science & Technology Commission (D171100003017002), and the National Science and Technology Major Project (2017ZX09304018).

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The study was approved by the ethics committees of Beijing Tiantan Hospital and all other research centres according to the principles expressed in the Declaration of Helsinki. Written informed consent was obtained from all patients or legally authorised representatives before entering into the study.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available to researchers on request for purposes of reproducing the results or replicating the procedure by directly contacting the corresponding author.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Anxin Wang <http://orcid.org/0000-0003-4351-2877>

Hongqiu Gu <http://orcid.org/0000-0003-1608-1856>

Hao Li <http://orcid.org/0000-0002-8591-4105>

Yongjun Wang <http://orcid.org/0000-0002-9976-2341>

REFERENCES

- Li S, Zhu L, Cheng X, *et al.* The significance of CO₂ combining power in predicting prognosis of patients with stage II and III colorectal cancer. *Biomark Med* 2019;13:1071–80.
- Hu J, Wang Y, Geng X, *et al.* Metabolic acidosis as a risk factor for the development of acute kidney injury and hospital mortality. *Exp Ther Med* 2017;13:2362–74.
- Ago T, Matsuo R, Hata J, *et al.* Insulin resistance and clinical outcomes after acute ischemic stroke. *Neurology* 2018;90:e1470–7.
- Mandel El, Forman JP, Curhan GC, *et al.* Plasma bicarbonate and odds of incident hypertension. *Am J Hypertens* 2013;26:1405–12.
- de Oliveira CMC, Vidal CLdaC, Cristino EF, *et al.* Metabolic acidosis and its association with nutritional status in hemodialysis. *J Bras Nefrol* 2015;37:458–66.
- Mahajan A, Simoni J, Sheather SJ, *et al.* Daily oral sodium bicarbonate preserves glomerular filtration rate by slowing its decline in early hypertensive nephropathy. *Kidney Int* 2010;78:303–9.
- Kendrick J, Shah P, Andrews E, *et al.* Effect of treatment of metabolic acidosis on vascular endothelial function in patients with CKD: a pilot randomized cross-over study. *Clin J Am Soc Nephrol* 2018;13:1463–70.
- Forlivesi S, Micheletti N, Tomelleri G, *et al.* Association of hyperglycemia, systolic and diastolic hypertension, and hyperthermia relative to baseline in the acute phase of stroke with poor outcome after intravenous thrombolysis. *Blood Coagul Fibrinolysis* 2018;29:167–71.
- Tiruvoipati R, Pilcher D, Botha J, *et al.* Association of hypercapnia and hypercapnic acidosis with clinical outcomes in mechanically ventilated patients with cerebral injury. *JAMA Neurol* 2018;75:818–26.
- Wang Y, Jing J, Meng X, *et al.* The third China national stroke registry (CNSR-III) for patients with acute ischaemic stroke or transient ischaemic attack: design, rationale and baseline patient characteristics. *Stroke Vasc Neurol* 2019;4:158–64.
- Adams HP, Bendixen BH, Kappelle LJ, *et al.* Classification of subtype of acute ischemic stroke. definitions for use in a multicenter clinical trial. TOAST. trial of ORG 10172 in acute stroke treatment. *Stroke* 1993;24:35–41.
- Thompson MP, Luo Z, Gardiner J, *et al.* Impact of missing stroke severity data on the accuracy of hospital ischemic stroke mortality profiling. *Circ Cardiovasc Qual Outcomes* 2018;11:e004951.
- Pottel H, Delanaye P, Schaeffner E, *et al.* Estimating glomerular filtration rate for the full age spectrum from serum creatinine and cystatin C. *Nephrol Dial Transplant* 2017;32:gfw425–507.
- Djamali A, Singh T, Melamed ML, *et al.* Metabolic acidosis 1 year following kidney transplantation and subsequent cardiovascular events and mortality: an observational cohort study. *Am J Kidney Dis* 2019;73:476–85.
- Dobre M, Yang W, Pan Q, *et al.* Persistent high serum bicarbonate and the risk of heart failure in patients with chronic kidney disease (CKD): a report from the chronic renal insufficiency cohort (CRIC) study. *J Am Heart Assoc* 2015;4. doi:10.1161/JAHA.114.001599. [Epub ahead of print: 20 Apr 2015].
- Kovesdy CP, Anderson JE, Kalantar-Zadeh K. Association of serum bicarbonate levels with mortality in patients with non-dialysis-dependent CKD. *Nephrol Dial Transplant* 2009;24:1232–7.
- Navaneethan SD, Schold JD, Arrigain S, *et al.* Serum bicarbonate and mortality in stage 3 and stage 4 chronic kidney disease. *Clin J Am Soc Nephrol* 2011;6:2395–402.
- Palmer BF. Evaluation and treatment of respiratory alkalosis. *Am J Kidney Dis* 2012;60:834–8.
- Li Y, Horiuchi T, Murata T, *et al.* Mechanism of alkalosis-induced constriction of rat cerebral penetrating arterioles. *Neurosci Res* 2011;70:98–103.
- Sun L, Zhang K, Li J, *et al.* An impairment of cortical GABAergic neurons is involved in alkalosis-induced brain dysfunctions. *Biochem Biophys Res Commun* 2012;419:627–31.
- Hamdi H, Hassanian-Moghaddam H, Hamdi A, *et al.* Acid-Base disturbances in acute poisoning and their association with survival. *J Crit Care* 2016;35:84–9.
- Kazmaier S, Weyland A, Buhre W, *et al.* Effects of respiratory alkalosis and acidosis on myocardial blood flow and metabolism in patients with coronary artery disease. *Anesthesiology* 1998;89:831–7.
- Al-Kindi SG, Sarode A, Zullo M, *et al.* Serum bicarbonate concentration and cause-specific mortality: the National health and nutrition examination survey 1999–2010. *Mayo Clin Proc* 2020;95:113–23.
- Raphael KL. Metabolic acidosis in CKD: core curriculum 2019. *Am J Kidney Dis* 2019;74:263–75.
- Kendrick JB, Zelnick L, Chonchol MB, *et al.* Serum bicarbonate is associated with heart failure in the multi-ethnic study of atherosclerosis. *Am J Nephrol* 2017;45:118–26.

SUPPLEMENTAL MATERIAL

Table S1. Comparison of baseline characteristics of participants and nonparticipants

Characteristics	Nonparticipants	Participants	<i>P</i> value
n	9531	5635	
Age, y	63.0(54.0-71.0)	62.0(54.0-70.0)	0.0048
Men, n (%)	3808(67.58)	6556(68.79)	0.1221
BMI, kg/m ²	24.24(22.49-26.29)	24.51(22.67-26.67)	<0.0001
Medical History, n (%)			
Hypertension	3536(62.75)	5958(62.51)	0.7689
Diabetes mellitus	1267(22.48)	2243(23.53)	0.1387
Dyslipidemia	390(6.92)	801(8.40)	0.0010
Stroke or TIA	1165(20.67)	2190(22.98)	0.0010
Atrial fibrillation/flutter	398(7.06)	621(6.52)	0.1932
Peripheral vascular disease	52(0.92)	66(0.69)	0.1188
Heart failure	30(3.77)	64(4.81)	0.2589
COPD	51(0.91)	81(0.85)	0.7236
Stroke type/Subtype, n (%)			
Ischemic stroke	5314(94.30)	8832(92.67)	0.0001
TIA	321(5.70)	699(7.33)	
TOAST, n (%)			
Large-artery atherosclerosis	1456(25.84)	2400(25.18)	0.0167
Cardioembolism	373(6.62)	544(5.71)	
Small-vessel occlusion	1199(21.28)	1966(20.63)	
Other determined etiology	56(0.99)	126(1.32)	
Undetermined etiology	2551(45.27)	4495(47.16)	
Current smoker, n (%)	1663(29.51)	3089(32.41)	0.0005
Medication in hospital, n (%)			

Cholesterol-lowering agents	5324(95.31)	9182(96.99)	<0.0001
Antihypertensive agents	2432(43.54)	4568(48.25)	<0.0001
Hypoglycemic agents	1343(24.04)	2449(25.87)	0.0126
Antiplatelet agents	5422(97.06)	9191(97.08)	0.9424
Anticoagulant agents	697(12.48)	849(8.97)	<0.0001
NIHSS score on admission	3(1-6)	3(1-6)	0.0010
Time from onset of symptoms to admission, h	13.00 (3.00-41.00)	15.00(3.00-45.00)	0.0213
Lipid level			
TC, mmol/L	3.97(3.31-4.69)	3.97(3.31-4.74)	0.7080
LDL, mmol/L	2.32(1.75-2.98)	2.30(1.71-2.97)	0.2338
HDL, mmol/L	0.94(0.78-1.13)	0.93(0.77-1.11)	0.0060
TG, mmol/L	1.35(1.02-1.85)	1.38(1.04-1.89)	0.0385
FBG, mmol/L	5.48(4.87-6.74)	5.56(4.90-6.99)	0.0028
eGFR, mL/min/1.73 m ²	92.73(81.15-101.75)	93.31(81.94-102.03)	0.1256
hs-CRP, mg/L	1.77(0.83-4.75)	1.78(0.81-4.71)	0.9442
Medication on discharge, n (%)			
Cholesterol-lowering agents	5080(90.33)	8571(92.06)	0.0003
Antihypertensive agents	2578(45.84)	4840(50.91)	<0.0001
Hypoglycemic agents	1248(22.19)	2298(24.17)	0.0054
Antiplatelet agents	5034(89.67)	8668(91.17)	0.0021
Anticoagulant agents	189(3.36)	260(2.73)	0.0284

Abbreviations: BMI=body mass index; COPD= chronic obstructive pulmonary disease; TIA=transient Ischemic Attack; TOAST= Trial of Org 10172 in Acute Stroke Treatment, NIHSS=The National Institutes of Health Stroke Scale; TC=total cholesterol; LDL=low-density lipoprotein cholesterol; HDL=high-density lipoprotein cholesterol; TG=triglycerides; FBG=fasting blood glucose; eGFR=estimated glomerular filtration rate; hs-CRP=high sensitivity C-reactive protein. Continuous variables are expressed as median with interquartile range. Categorical variables are expressed as frequency with percentage.

Table S2. Subgroup analyses for the associations of CO₂CP with all-cause death and poor functional outcomes.

Outcomes	Quintiles of CO ₂ CP, mmol/L					<i>P</i> _{interaction}
	Q1	Q2	Q3	Q4	Q5	
Age						
At 3 months						
Age<65 years						
Death	0.81(0.26-2.49)	0.48(0.15-1.52)	0.38(0.10-1.38)	Reference	1.21(0.41-3.58)	0.1507
mRS score 3-6	1.36(0.97-1.91)	1.16(0.83-1.62)	1.05(0.75-1.47)	Reference	1.02(0.72-1.44)	0.7900
mRS score 2-6	1.20(0.93-1.55)	1.11(0.87-1.42)	0.97(0.76-1.24)	Reference	1.11(0.86-1.42)	0.5358
Age≥ 65 years						
Death	4.53(1.87-10.96)	2.49(1.00-6.18)	2.13(0.84-5.40)	Reference	1.61(0.63-4.12)	
mRS score 3-6	1.65(1.20-2.27)	1.32(0.96-1.80)	1.38(1.00-1.90)	Reference	0.96(0.70-1.33)	
mRS score 2-6	1.26(0.97-1.63)	1.04(0.80-1.33)	1.15(0.89-1.48)	Reference	0.95(0.74-1.22)	
At 1 year						
Age<65 years						
Death	0.91(0.41-2.03)	0.62(0.28-1.42)	1.06(0.50-2.26)	Reference	1.27(0.60-2.69)	0.0937

mRS score 3-6	1.38(0.96-1.98)	1.08(0.75-1.55)	1.12(0.79-1.60)	Reference	1.22(0.85-1.76)	0.5234
mRS score 2-6	1.33(1.02-1.73)	1.12(0.87-1.44)	1.16(0.91-1.49)	Reference	1.12(0.87-1.45)	0.8738
Age ≥ 65 years						
Death	1.88(1.42-2.53)	1.46(1.10-1.94)	1.42(1.06-1.89)	Reference	1.02(0.77-1.37)	
mRS score 3-6	1.24(0.91-1.68)	0.94(0.69-1.27)	1.08(0.79-1.46)	Reference	0.88(0.65-1.19)	
mRS score 2-6	1.33(1.02-1.72)	1.14(0.89-1.46)	1.12(0.87-1.45)	Reference	0.98(0.77-1.27)	
Gender						
At 3 months						
Male						
Death	2.11(0.85-5.23)	2.53(1.03-6.16)	1.22(0.47-3.18)	Reference	1.95(0.77-4.89)	0.4843
mRS score 3-6	1.42(1.07-1.89)	1.24(0.93-1.65)	1.33(1.01-1.76)	Reference	0.94(0.70-1.26)	0.4217
mRS score 2-6	1.16(0.93-1.45)	1.05(0.85-1.31)	1.02(0.82-1.26)	Reference	1.04(0.84-1.29)	0.9366
Female						
Death	2.68(1.03-6.96)	1.19(0.62-2.28)	1.46(0.77-2.78)	Reference	0.59(0.27-1.29)	

mRS score 3-6	2.12(1.43-3.14)	1.38(0.94-2.02)	1.10(0.73-1.65)	Reference	1.08(0.72-1.60)	
mRS score 2-6	1.62(1.17-2.23)	1.23(0.91-1.67)	1.27(0.93-1.75)	Reference	1.06(0.76-1.46)	
At 1 year						
Male						
Death	1.31(0.75-2.28)	1.53(0.89-2.64)	1.28(0.74-2.22)	Reference	1.47(0.85-2.55)	0.2021
mRS score 3-6	1.28(0.96-1.70)	1.09(0.82-1.46)	1.14(0.86-1.51)	Reference	1.07(0.80-1.44)	0.8290
mRS score 2-6	1.27(1.02-1.59)	1.15(0.92-1.43)	1.10(0.89-1.36)	Reference	1.09(0.88-1.36)	0.8300
Female						
Death	2.12(1.07-4.17)	1.24(0.62-2.49)	1.53(0.77-3.03)	Reference	0.60(0.26-1.36)	
mRS score 3-6	1.46(0.99-2.16)	0.95(0.65-1.39)	1.09(0.74-1.61)	Reference	0.89(0.60-1.32)	
mRS score 2-6	1.61(1.17-2.22)	1.20(0.89-1.63)	1.38(1.01-1.90)	Reference	1.02(0.74-1.40)	
Stroke subtypes						
At 3 months						
Ischemic stroke						

Death	2.38(1.26-4.51)	1.57(0.82-3.01)	1.16(0.58-2.31)	Reference	1.28(0.64-2.54)	0.7622
mRS score 3-6	1.56(1.24-1.97)	1.26(1.00-1.58)	1.24(0.98-1.56)	Reference	0.95(0.75-1.20)	0.7137
mRS score 2-6	1.23(1.03-1.48)	1.08(0.91-1.29)	1.07(0.89-1.27)	Reference	1.01(0.85-1.21)	0.8722
TIA						
Death				Reference		
mRS score 3-6	0.98(0.06-16.29)	1.35(0.12-15.39)	1.66(0.16-17.75)	Reference	0.98(0.08-11.71)	
mRS score 2-6	2.52(0.57-11.09)	1.85(0.45-7.67)	2.13(0.55-8.30)	Reference	1.50(0.37-6.03)	
At 1 year						
Ischemic stroke						
Death	1.56(1.02-2.38)	1.33(0.87-2.03)	1.32(0.86-2.01)	Reference	1.05(0.68-1.65)	0.9907
mRS score 3-6	1.34(1.07-1.69)	0.99(0.79-1.25)	1.11(0.88-1.40)	Reference	0.98(0.78-1.23)	0.2221
mRS score 2-6	1.35(1.12-1.62)	1.13(0.95-1.36)	1.16(0.97-1.38)	Reference	1.07(0.89-1.28)	0.8889
TIA						
Death	7.63(0.02-3626.6)	1.38(0.02-84.45)	30.49	Reference	2.19(0.01-610)	

mRS score 3-6	0.62(0.06-6.61)	1.55(0.23-10.46)	1.61(0.20-12.97)	Reference	1.01(0.11-9.03)	
mRS score 2-6	1.57(0.44-5.60)	1.09(0.34-3.46)	2.35(0.77-7.19)	Reference	0.75(0.24-2.36)	
Smoking						
Current smoking						
Death	1.69(0.30-9.54)	3.34(0.71-15.67)	0.99(0.18-5.57)	Reference	1.54(0.30-7.82)	0.1856
mRS score 3-6	1.83(1.17-2.85)	1.54(0.98-2.44)	1.51(0.97-2.36)	Reference	1.38(0.87-2.21)	0.7411
mRS score 2-6	1.39(1.00-1.93)	1.13(0.80-1.58)	1.02(0.74-1.42)	Reference	1.28(0.91-1.80)	0.9321
No smoking						
Death	2.59 (1.28-5.23)	1.31(0.62-2.79)	1.03(0.47-2.25)	Reference	1.15(0.53-2.52)	
mRS score 3-6	1.60(1.22-2.10)	1.23(0.94-1.60)	1.19(0.91-1.56)	Reference	0.91(0.69-1.20)	
mRS score 2-6	1.25(1.01-1.56)	1.10(0.89-1.35)	1.08(0.88-1.34)	Reference	0.96(0.76-1.18)	
At 1 year						
Current smoking						
Death	2.17(0.77-6.11)	2.44(0.86-6.93)	2.03(0.73-5.68)	Reference	1.87(0.63-5.56)	0.8225

mRS score 3-6	1.47(0.95-2.28)	0.99(0.62-1.58)	1.27(0.81-1.97)	Reference	1.07(0.67-1.72)	0.6878
mRS score 2-6	1.56(1.12-2.18)	1.24(0.88-1.75)	1.33(0.96-1.85)	Reference	1.36(0.96-1.91)	0.4133
No smoking						
Death	1.38(0.87-2.21)	1.17(0.74-1.86)	1.16(0.73-1.84)	Reference	0.88(0.53-1.44)	
mRS score 3-6	1.28(0.97-1.68)	1.02(0.79-1.33)	1.07(0.82-1.40)	Reference	0.97(0.74-1.26)	
mRS score 2-6	1.30(1.04-1.61)	1.13(0.92-1.38)	1.11(0.90-1.37)	Reference	0.96(0.78-1.19)	
NIHSS						
At 3 months						
NIHSS<4						
Death	3.87(1.40-10.68)	1.61(0.54-4.77)	1.30(0.41-4.18)	Reference	0.92(0.29-2.86)	0.4801
mRS score 3-6	1.45(1.01-2.10)	1.04(0.72-1.49)	1.10(0.76-1.58)	Reference	0.78(0.54-1.14)	0.4643
mRS score 2-6	1.11(0.86-1.44)	0.98(0.77-1.24)	0.93(0.73-1.18)	Reference	0.93(0.73-1.18)	0.1799
NIHSS ≥ 4						
Death	1.36(0.56-3.26)	1.42(0.60-3.37)	1.03(0.41-2.59)	Reference	1.54(0.62-3.83)	

mRS score 3-6	1.61(1.20-2.17)	1.37(1.02-1.84)	1.48(1.10-1.99)	Reference	1.10(0.81-1.50)	
mRS score 2-6	1.42(1.09-1.85)	1.18(0.91-1.53)	1.33(1.03-1.73)	Reference	1.09(0.84-1.43)	
At 1 year						
NIHSS<4						
Death	2.12(1.11-4.04)	1.50(0.80-2.85)	1.35(0.69-2.64)	Reference	0.88(0.43-1.78)	0.3072
mRS score 3-6	1.29(0.91-1.84)	0.95(0.67-1.35)	1.08(0.77-1.53)	Reference	0.94(0.66-1.33)	0.9444
mRS score 2-6	1.43(1.11-1.84)	1.21(0.96-1.54)	1.13(0.89-1.44)	Reference	1.06(0.83-1.35)	0.3117
NIHSS ≥ 4						
Death	1.14(0.65-2.01)	1.05(0.60-1.86)	1.23(0.70-2.15)	Reference	1.26(0.71-2.26)	
mRS score 3-6	1.35(0.99-1.84)	1.01(0.74-1.39)	1.21(0.88-1.65)	Reference	1.04(0.76-1.44)	
mRS score 2-6	1.36(1.03-1.77)	1.04(0.80-1.36)	1.30(0.99-1.69)	Reference	1.06(0.81-1.39)	

Abbreviations: CO₂CP=carbon dioxide combining power; mRS=modified Rankin Scale; NIHSS=The National Institutes of Health Stroke Scale; TIA=transient ischemic attack

Adjusted for age, gender, body mass index, history of dyslipidemia, stroke or transient ischemic stroke, atrial fibrillation/flutter, stroke type, current smoke, antihypertensive agents, anticoagulant agents, NIHSS, high-density lipoprotein cholesterol, history of COPD, time from onset of symptoms to admission, and high sensitivity C-reactive protein on admission.

Table S3. Comparison of baseline characteristics of patients stratified by CO₂CP levels.

Characteristics	CO ₂ CP levels			P value
	<23 mmol/L	23-29 mmol/L	≥29 mmol/L	
n				
Age, y	62(53-70)	62(54-70)	63.5(55-70)	0.0122
Men, n (%)	1832(69.53)	4109(68.64)	615(67.58)	0.5112
BMI, kg/m ²	24.62(22.86-26.70)	24.49(22.66-26.64)	24.61(22.58-26.73)	0.2998
Medical History, n (%)				
Hypertension	1632(61.94)	3751(62.66)	575(63.19)	0.7276
Diabetes mellitus	590(22.39)	1440(24.06)	213(23.41)	0.2431
Dyslipidemia	187(7.10)	549(9.17)	65(7.14)	0.0021
Stroke or TIA	567(21.52)	1388(23.19)	235(25.82)	0.0237
Atrial fibrillation/flutter	197(7.48)	381(6.36)	43(4.73)	0.0111
Peripheral vascular disease	18(0.68)	40(0.67)	8(0.88)	0.7728
Heart failure	34(4.14)	21(5.72)	9(6.29)	0.3401
COPD	22 (0.83)	49 (0.82)	10(1.10)	0.6886
Stroke type/Subtype, n (%)				
Ischemic stroke	5527(92.33)	2469(93.70)	836(91.87)	0.0503
TIA	459(7.67)	166(6.30)	74(8.13)	

TOAST, n (%)				
Large-artery atherosclerosis	1486(24.82)	687(26.07)	227(24.95)	0.2416
Cardioembolism	336(5.61)	165(6.26)	43(4.73)	
Small-vessel occlusion	1269(21.20)	499(18.94)	198(21.76)	
Other determined etiology	80(1.34)	37(1.40)	9(0.99)	
Undetermined etiology	2815(47.03)	1247(46.32)	433(47.58)	
Current smoker, n (%)	1895(31.66)	921(34.95)	273(30.00)	0.0006
Medication in hospital, n (%)				
Cholesterol-lowering agents	5768(96.99)	2538(96.98)	876(97.01)	0.9990
Antihypertensive agents	2823(48.25)	1320(50.44)	425(47.07)	0.0305
Hypoglycemic agents	1559(26.21)	661(25.26)	229(25.36)	0.6057
Antiplatelet agents	5767(96.97)	2552(97.52)	872(96.57)	0.2420
Anticoagulant agents	491(8.26)	291(11.12)	67(7.42)	<0.0001
NIHSS score on admission	3(1-6)	3(1-6)	3(1-5)	<0.0001
Time from onset of symptoms to admission, h	13.00 (3.00-34.00)	15.00(3.00-46.00)	18.00(4.00-48.00)	0.0009
Lipid level				
TC, mmol/L	3.98(3.32-4.73)	3.94(3.31-4.75)	3.97(3.31-4.77)	0.8104
LDL, mmol/L	2.32(1.71-2.97)	2.27(1.70-2.97)	2.27(1.70-2.97)	0.5744

HDL, mmol/L	0.93(0.77-1.11)	0.91(0.77-1.12)	0.95(0.79-1.09)	0.0504
TG, mmol/L	1.39(1.06-1.91)	1.38(1.03-1.89)	1.37(1.05-1.86)	0.7563
FBG, mmol/L	5.58(4.89-6.95)	5.56(4.90-6.97)	5.52(4.90-7.10)	0.9095
eGFR, mL/min/1.73 m ²	92.83(80.74-102.55)	93.72(82.37-101.96)	92.83(82.72-101.01)	0.3129
hs-CRP, mg/L	1.97(0.87-5.47)	1.72(0.80-4.53)	1.62(0.79-4.21)	0.0011
Medication on discharge, n (%)				
Cholesterol-lowering agents	2440 (93.99)	5472(91.60)	839(92.30)	0.0861
Antihypertensive agents	1387(52.86)	3003(50.27)	450(49.50)	0.0582
Hypoglycemic agents	637(24.48)	1446(24.20)	215(23.65)	0.9264
Antiplatelet agents	2400(91.46)	5434(900.96)	834(91.75)	0.6113
Anticoagulant agents	86(3.28)	151(2.53)	23(2.53)	0.1345

Abbreviations: BMI=body mass index; CO₂CP=carbon dioxide combining power; COPD= chronic obstructive pulmonary disease; TIA=transient Ischemic Attack; TOAST=Trials of Org 10172 in Acute Stroke Treatment; NIHSS=The National Institutes of Health Stroke Scale; TC=total cholesterol; LDL=low-density lipoprotein cholesterol; HDL=high-density lipoprotein cholesterol; TG=triglycerides; FBG=fasting blood glucose; eGFR=estimated glomerular filtration rate; hs-CRP=high sensitivity C-reactive protein. Continuous variables are expressed as median with interquartile range. Categorical variables are expressed as frequency with percentage.

Table S4. Reclassification and discrimination statistics for outcomes at 3 months and 1 year by CO₂CP

	C statistics		IDI		NRI (continuous)		NRI (categorical) ^a	
	Estimate (95% CI)	<i>P</i> value	Estimate (95% CI), %	<i>P</i> value	Estimate (95% CI), %	<i>P</i> value	Estimate (95% CI), %	<i>P</i> value
Outcomes at 3 months								
Death								
Conventional model ^b	0.841 (0.805-0.876)		Reference		Reference		Reference	
Conventional model + CO ₂ CP	0.843 (0.807-0.878)	0.6886	0.50 (0.06-0.94)	0.0254	19.43 (1.91-36.95)	0.0303	2.34 (-1.76-6.43)	0.2657
mRS score 3-6								
Conventional model ^b	0.809 (0.796-0.822)		Reference		Reference		Reference	
Conventional model + CO ₂ CP	0.810 (0.797-0.823)	0.3550	0.20 (0.07-0.34)	0.0024	10.72 (4.86-16.58)	0.0003	0.02 (-1.59-1.23)	0.8004
mRS score 2-6								
Conventional model ^b	0.783 (0.773-0.795)		Reference		Reference		Reference	
Conventional model + CO ₂ CP	0.784 (0.774-0.795)	0.7612	0.03 (-0.01-0.01)	0.1497	4.82 (0.25-9.39)	0.0388	0.01 (-0.1-0.05)	0.6700
Outcomes at 1 year								
Death								
Conventional model ^b	0.795 (0.768-0.823)		Reference		Reference		Reference	
Conventional model + CO ₂ CP	0.798 (0.770-0.825)	0.3269	0.18 (-0.01-0.41)	0.1431	11.80 (-0.01-23.67)	0.0518	1.03 (-0.13-3.35)	0.3868
mRS score 3-6								

Conventional model ^b	0.799 (0.786-0.813)		Reference		Reference		Reference	
Conventional model + CO ₂ CP	0.801 (0.787-0.814)	0.1937	0.11 (0.01-0.23)	0.0353	6.49 (0.51-12.46)	0.0334	0.18 (-1.05-1.40)	0.7790
mRS score 2-6								
Conventional model ^b	0.761 (0.750-0.772)		Reference		Reference		Reference	
Conventional model + CO ₂ CP	0.762 (0.751-0.773)	0.1822	0.07 (0.01-0.14)	0.0383	4.74 (0.04-9.43)	0.0483	0.03 (-0.12-0.06)	0.5369

Abbreviations: CI=confidence interval; CO₂CP=carbon dioxide combining power; IDI=integrated discrimination improvement; NRI=net reclassification index; mRS=modified Rankin Scale.

Patients were divided into 5 risk categories by CO₂CP quintiles.

Conventional model: adjusted for age, gender, body mass index, history of dyslipidemia, stroke or transient ischemic stroke, atrial fibrillation/flutter, stroke type, current smoke, antihypertensive agents, anticoagulant agents, NIHSS, high-density lipoprotein cholesterol, history of COPD, time from onset of symptoms to admission, and high sensitivity C-reactive protein on admission.